



UNIVERSITI PUTRA MALAYSIA

**ENDOPHYTIC MICROORGANISMS FOR IMPROVEMENT OF
BANANA VIGOUR AND TOLERANCE TO FUSARIUM WILT**

ADELINE TING SU YIEW.

FP 2005 9

**ENDOPHYTIC MICROORGANISMS FOR IMPROVEMENT OF BANANA
VIGOUR AND TOLERANCE TO FUSARIUM WILT**

By

ADELINE TING SU YIEN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of Doctor of Philosophy**

October 2005



DEDICATIONS

**I dedicate the fruits of my labour
to my loving late Mum,
to my understanding family,
and my affectionate husband Steve.
Thank you for believing in me.**

Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Doctor of Philosophy

**ENDOPHYTIC MICROORGANISMS FOR IMPROVEMENT OF BANANA
VIGOUR AND TOLERANCE TO FUSARIUM WILT**

By

ADELINE TING SU YIEN

October 2005

Chairperson: Professor Sariah Meon, PhD

Faculty : Agriculture

This study explored the potential of endophytic microorganisms (EMs) isolated from wild bananas as biocontrol agents (BCAs) against Fusarium wilt caused by *Fusarium oxysporum* f. sp. *cubense* race 4 (FocR4) in susceptible banana ramets (Berangan cv. Intan).

The fungal (*Fusarium oxysporum* (UPM31P1)) and bacterial endophytes (*Serratia marcescens* (UPM39B3)) were established as effective BCAs; improving growth and vigour and enhancing tolerance of susceptible ramets to Fusarium wilt. They were antagonistic towards FocR4, with Percentage Inhibition of Radial Growth (PIRG) values of 65% and 63%, respectively. Inhibitory substances were produced in the form of volatile as well as non-volatile substances. The endophytes were also able to colonize the host tissues,

including the roots and corms in artificial inoculation under laboratory conditions. The association of *F. oxysporum* (UPM31P1) and *S. marcescens* (UPM39B3) with the host plants resulted in enhanced vegetative growth as shown by the increase in height, pseudostem diameter, root mass and total number of leaves ramet¹. Endophytes acted as elicitors in the production of inducible compounds associated with induced resistance (peroxidase, polyphenoloxidase, phenylalanine ammonia lyase, total soluble phenol and lignin content). The robust growth and occurrence of induced resistance subsequently enhanced tolerance of the ramets to Fusarium wilt based on parameters such as delay in onset of symptoms, lower percentages in disease incidence, disease severity, and epidemic rate. Inoculation with *F. oxysporum* singly was most effective followed by inoculation with mixture with *S. marcescens* (UPM31P1+UPM39B3). However, the survival and subsequent biocontrol efficacy of *F. oxysporum* (UPM31P1) and *S. marcescens* (UPM39B3) might be vulnerable to the changing soil and environmental conditions. Populations of endophytes were not sustained over time. Therefore, further studies regarding formulation and application frequency and techniques, are essential to maximize the potential of *F. oxysporum* (UPM31P1) and *S. marcescens* (UPM39B3) as BCAs against Fusarium wilt of banana.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**MIKROORGANISMA ENDOFIT UNTUK MENINGKATKAN TUMBESARAN
DAN TOLERANSI TANAMAN PISANG TERHADAP
PENYAKIT LAYU FUSARIUM**

Oleh

ADELINE TING SU YIEN

Oktober 2005

Pengerusi : Profesor Sariah Meon, PhD

Fakulti : Pertanian

Kajian ini mengeksplorasi potensi mikroorganisma endofit (EMs) yang diasingkan dari pokok pisang liar sebagai agen kawalan biologi (BCAs) untuk menangani penyakit layu Fusarium yang disebabkan oleh *Fusarium oxysporum* f. sp. *cubense* ras 4 (FocR4) pada ramet pisang (Berangan cv. Intan).

Isolat kulat *Fusarium oxysporum* (UPM31P1) dan bakteria *Serratia marcescens* (UPM39B3) dikenalpasti sebagai agen kawalan biologi yang efektif; meningkatkan tumbesaran dan toleransi ramet pisang terhadap layu Fusarium. Kedua-dua isolat ini adalah antagonistik terhadap FocR4, masing-masing dengan peratus perencatan pertumbuhan miselium (PIRG) 65% dan 63%. Bahan perencatan dikeluarkan dalam bentuk bahan mudah meruap dan tidak mudah meruap. Kedua-dua endofit ini juga berupaya menjajah tisu perumah, termasuk

tisu akar dan umbisi, melalui kaedah inokulasi tiruan. Gabungan *F. oxysporum* (UPM31P1) dan *S. marcescens* (UPM39B3) dengan perumah menghasilkan peningkatan pada tumbesaran vegetatif seperti yang ditunjukkan oleh peningkatan dalam ketinggian, lilitan pseudostem, massa akar dan jumlah bilangan daun ramet¹. Endofit bertindak sebagai elisitor dalam pengeluaran kompaun teraruh berkaitan dengan keresistanan teraruh (peroksidase, polifenoloksidase, fenelalanin ammonia liase, fenol terlarut dan kandungan lignin). Peningkatan tumbesaran dan pengeluaran kompaun teraruh meningkatkan ketahanan ramet pisang terhadap layu Fusarium berdasarkan kepada parameter seperti pelambatan kemunculan simptom, pengurangan peratus insiden dan keterukan penyakit dan kadar epidemik. Penginokulatan dengan *F. oxysporum* (UPM31P1) sahaja adalah paling efektif, diikuti dengan inokulasi secara kombinasi dengan *S. marcescens* (UPM31P1+UPM39B3). Namun demikian, kemandirian dan keberkesanan kawalan biologi *F. oxysporum* (UPM31P1) dan *S. marcescens* (UPM39B3) mungkin dipengaruhi oleh keadaan tanah dan alam sekitar yang tidak menentu. Populasi kedua-dua isolat endofit juga menurun mengikut masa. Oleh itu, kajian seterusnya mengenai formulasi dan frekuensi dan teknik aplikasi adalah penting untuk memaksimumkan potensi *F. oxysporum* (UPM31P1) dan *S. marcescens* (UPM39B3) sebagai agen kawalan biologi untuk layu Fusarium pada pisang.

ACKNOWLEDGEMENTS

I thank God for His grace and blessings, in the many moments of my life. Praise be to God for His sovereign love, mercy and faithfulness.

I extend my gratitude to Professor Dr. Sariah Meon, for her constant guidance and constructive criticisms throughout the supervision of the project. Also for her kind and sharing gesture for the many opportunities for scientific interaction availed to me. My heartfelt appreciation as well to Associate Professor Dr. Jugah Kadir, Professor Dr. Son Radu and Dr. Gurmit Singh, for their guidance and suggestions to continuously improve the project. Truly, I am exceedingly grateful for their commitments to this project.

My sincerest appreciation as well to the Malaysian Ministry of Science, Technology and Innovation (MOSTI) for the National Science Fellowship (NSF) awarded. I am also most grateful to the assistance provided by Madam Ho Yuk Wah and her team in United Plantations Bhd (Teluk Intan), not just during the course of my field-trial but the many times involved in preparations of plant materials. Many thanks as well to the staffs and officers from the Plant Pathology Laboratory and Plant Protection Department for their kind assistance.



And, finally, to my family and husband, I thank you for your love, patience, understanding and support, which has kept me accompanied through the high and low points throughout the period of my study. Especially to Steve, thank you for absorbing the weight of my anxieties so I shall not stand alone.

I certify that an Examination Committee met on 13th October 2005 to conduct the final examination of Adeline Ting Su Yien on her Doctor of Philosophy thesis entitled "Endophytic Microorganisms for Improvement of Banana Vigour and Tolerance to Fusarium Wilt" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Tan Yee How, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Zulkifli Shamsuddin, PhD


Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Suhaimi Napis, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Baharuddin Salleh, PhD

Professor
School of Biological Sciences
Universiti Sains Malaysia
(External Examiner)



HASANAH MOHD. GHAZALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **19 JAN 2006**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Sariah Meon, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Jugah Kadir, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Son Radu, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Gurmit Singh, PhD

Director of Research Department
United Plantations Bhd
Jenderata Estate, Teluk Intan
(Member)



AINI IDERIS, PhD

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 07 FEB 2006

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



ADELINE TING SU YIEN

Date: 12.12.2005.

TABLE OF CONTENTS

DEDICATIONS	Page ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF PLATES	xxi
LIST OF ABBREVIATIONS	xxiv

CHAPTER

1	INTRODUCTION	1.1
	1.1 Objectives of Study	1.4
	1.2 Hypothesis	1.4
 2	 LITERATURE REVIEW	 2.1
	2.1 The Banana Profile	2.1
	2.2 Threat of <i>Fusarium</i> Wilt to the Banana Industry	2.3
	2.3 The Pathogen: <i>Fusarium oxysporum</i> f. sp. <i>cubense</i>	2.5
	2.4 Disease Incidence: Epidemiology and Predisposing Factors	2.7
	2.5 Control Measures	2.9
	2.5.1 Conventional Practices	2.9
	2.5.2 Plant-Breeding and Transgenic Approach	2.10
	2.5.3 Biological Management	2.14
	2.6 Endophytic Microorganisms	2.15
	2.6.1 Definition, Origin and Host Distribution	2.15
	2.6.2 Endophytic Microorganisms and Host Plant Interaction	2.18
	2.6.3 Fungal Endophytes	2.20
	2.6.4 Bacterial Endophytes	2.22
	2.6.5 Endophytic Microorganisms and their Potential as Biological Control Agents (BCAs)	2.23
	2.7 Introduction of Endophytic Microorganisms to Host Plants - Inoculation Techniques	2.26



2.8	Host Defense Mechanism-Induced Resistance	2.27
2.9	Molecular Markers in Microbial Identification	2.34
2.9.1	Conventional Fungal Typing Techniques	2.35
2.9.2	Conventional Bacterial Typing Techniques	2.36
2.9.3	Microbial Identification Using Molecular Markers	2.36
2.9.4	The RAPD-PCR Technique	2.38
3	ISOLATION AND CHARACTERIZATION OF ENDOPHYTIC MICROORGANISMS (EMs) FROM WILD BANANA	3.1
3.1	Introduction	3.1
3.2	Materials and Methods	3.2
3.2.1	Detection of Endophytic Microorganisms (EMs) in Host Tissues	3.2
3.2.2	Isolation of Endophytic Microorganisms (EMs)	3.4
3.2.3	Screening for Antagonistic Activity	3.5
3.2.4	Classification of Endophytic Microorganisms (EMs) Based on DNA Polymorphism	3.11
3.2.5	Identification of Endophytic Microorganisms (EMs)	3.17
3.3	Results and Discussion	3.21
3.3.1	Collection of Endophytic Microorganisms (EMs) from Wild Banana	3.19
3.3.2	Detection of Endophytic Microorganisms (EMs) in Host Tissues	3.20
3.3.3	Isolation of Endophytic Microorganisms(EMs)	3.20
3.3.4	Establishment of Antagonistic Activity	3.25
3.3.5	DNA Polymorphism of Fungal and Bacterial Endophytes	3.31
3.3.6	Identification of Biological Control Agent (BCA) Candidates	3.38
4	THE EFFECT OF SELECTED ENDOPHYTIC MICROORGANISMS (EMs) ON PRODUCTION OF INDUCIBLE COMPOUNDS AND GROWTH ASSOCIATED WITH INDUCED RESISTANCE IN BANANA RAMETS	4.1
4.1	Introduction	4.1
4.2	Materials and Methods	4.2
4.2.1	Experimental Design	4.2
4.2.2	Inoculum Preparation	4.2
4.2.3	Ramet Preparation and Inoculation	4.3
4.2.4	Disease Assessment	4.4

4.2.5	Effect of Endophytic Microorganisms (EMs) on Production of Inducible Compounds Associated with Induced Resistance	4.7
4.2.6	Effect of Endophytic Microorganisms (EMs) on Plant Vigour	4.11
4.2.7	Colonization and Establishment of Endophytic Microorganisms (EMs) in Host Tissues	4.11
4.3	Results and Discussion	4.15
4.3.1	Disease Assessment	4.15
4.3.2	Effect of Endophytic Microorganisms (EMs) on Production of Inducible Compounds Associated with Induced Resistance	4.16
4.3.3	Effect of Endophytic Microorganisms (EMs) on Plant Vigour	4.26
4.3.4	Colonization and Establishment of Endophytic Microorganisms (EMs) in Banana (Berangan cv. Intan)	4.29

5 THE EFFICACY OF ENDOPHYTIC MICROORGANISMS (EMs) IN SUPPRESSING FUSARIUM WILT INCIDENCE IN BANANAS

5.1	Introduction	5.1
5.2	Materials and Methods	5.1
5.2.1	Experimental Design	5.1
5.2.2	Inoculum Preparation	5.2
5.2.3	Ramet Preparation and Inoculation	5.3
5.2.4	Parameters for Disease Suppression Evaluation	5.3
5.2.5	Colonization and Establishment of Endophytic Microorganisms (EMs) in Host Tissues	5.8
5.2.6	Repeat Assessment on the Efficacy of Selected Endophytic Microorganisms (EMs) as Potential Biological Control Agents (BCAs)	5.9
5.3	Results and Discussion	5.10
5.3.1	Disease Assessment	5.10
5.3.2	Histopathological Observations of Healthy and Infected Banana Roots	5.19
5.3.3	Effect of Endophytic Microorganisms (EMs) on the Production of Inducible Compounds Associated with Induced Resistance	5.24
5.3.4	Effect of Endophytic Microorganisms (EMs) on Plant Vigour	5.35

5.3.5	Colonization of Endophytic Microorganisms (EMs) in Host Tissues	5.40
5.3.6	Repeat Disease Assessment on the Efficacy of Selected Endophytic Microorganisms (EMs) as Potential Biological Control Agents (BCAs)	5.45
6	FIELD EVALUATION ON THE EFFICACY OF UPM31P1 AND UPM39B3 FOR THE CONTROL OF FUSARIUM WILT	6.1
6.1	Introduction	6.1
6.2	Materials and Methods	6.1
6.2.1	Experimental Design	6.1
6.2.2	Mass Production of Inoculum	6.2
6.2.3	Ramet Preparation and Inoculation	6.4
6.2.4	Assessment of Biocontrol Efficacy	6.6
6.3	Results and Discussion	6.7
6.3.1	Development of Disease Symptoms	6.7
6.3.2	Effect of Endophytic Microorganisms (EMs) on Vegetative Growth	6.15
6.3.3	Inoculum Potential of Endophytes in Banana Plants	6.17
7	GENERAL DISCUSSION AND CONCLUSION	7.1
	REFERENCES	R.1
	APPENDICES	A.1
	BIODATA OF THE AUTHOR	B.1



LIST OF TABLES



Table	Page
3.1 Percentage of Inhibition of Radial Growth values (PIRG values, %) recorded by endophytic isolates against FocR4 in double plate (volatile) and poison-food test (non-volatile)	3.30
3.2 Cultural (on PDA) and morphological characteristics of UPM31F4 and UPM31P1	3.39
4.1 Production of inducible compounds associated with induced resistance in root tissues of banana ramets (Berangan cv. Intan) inoculated with endophytes. (Values are means of 20 replicates)	4.18
4.2 Production of inducible compounds associated with induced resistance in leaf tissues of banana ramets (Berangan cv. Intan) inoculated with endophytes. (Values are means of 20 replicates)	4.19
4.3 Vegetative growth (increase in height, pseudostem diameter, root mass, total number of leaves ramet ⁻¹) of ramets inoculated with endophytic isolates compared to positive (SDW) and negative (FocR4) control. (Values are means of 20 replicates)	4.27
4.4 Frequency (%) of isolation (recovery) of endophytes from root, corm, pseudostem and leaf tissue according to time	4.30
5.1 Different treatments of single or mixtures of endophytes on banana ramets (Berangan cv. Intan) for disease assessment	5.2
5.2 Different treatments of single or mixture of endophytes on banana ramets (Berangan cv. Intan) for repeat disease assessment trial	5.9
5.3 Values of Area Under Disease Progress Curve (AUDPC, units days after FocR4 challenged inoculation ⁻¹), % in disease reduction (DR%) and epidemic rate (r_m , units days ⁻¹) in banana ramets (Berangan cv. Intan) pre-inoculated with endophytes and challenged with FocR4	5.15
5.4 Production of inducible compounds associated with induced resistance in root tissues of banana ramets (Berangan cv. Intan) pre-inoculated with endophytes and challenged with FocR4 inoculation. (Values are means of 56 replicates)	5.26



5.5	Production of inducible compounds associated with induced resistance in leaf tissues of banana ramets (Berangan cv. Intan) pre-inoculated with endophytes and challenged with FocR4 inoculation. (Values are means of 56 replicates)	5.27
5.6	Vegetative growth (increase in height, pseudostem diameter, root mass, total number of leaves ramet ¹) of banana ramets (Berangan cv. Intan) pre-inoculated with endophytes and challenged with FocR4 inoculation. (Values are means of 56 replicates)	5.36
5.7	Population of endophytes recovered from different tissues of FocR4 infected and uninfected ramets pre-inoculated with endophytes. (Values are means of 56 replicates)	5.43
6.1	Time-frame of inoculation and population estimation periods of ramets pre-inoculated with UPM31P1 and UPM39B3	6.6

LIST OF FIGURES

Figure	Page
3.1 Total endophytic isolates recovered from wild banana roots on Fusarium Selective Media (FSM), Potato Dextrose Agar (PDA) and Nutrient Agar (NA), expressed as percentage (%) of total number of endophytes isolated	3.23
3.2 Total <i>Fusarium</i> spp. isolated from wild banana roots on Fusarium Selective Media (FSM), expressed as percentage (%) of total number of fungal endophytes isolated	3.23
3.3 Percentage of bacterial endophytes with antagonistic potential against FocR4 based on their respective PIRG values. (PIRG values are means of three replicates)	3.27
3.4 Percentage of fungal endophytes with antagonistic potential against FocR4 based on their respective PIRG values. (PIRG values are means of three replicates)	3.27
3.5 Dendrogram of fungal endophytes generated using OPC 11, 14 and 15. Cluster I: pathogenic isolates; cluster II to IX: nonpathogenic isolates. (Fungal endophytes selected as potential BCAs are isolates number 49 and 50 in cluster III and VII, respectively (▲✓))	3.33
3.6 Dendrogram of bacterial endophytes generated using OPC 1, 8 and 11. (Bacterial endophytes selected as potential BCAs are isolates number 48, 50 and 70 in cluster IV, V and VI, respectively (▲✓))	3.37
4.1 Population of UPM31F4 isolated from root, corm, pseudostem and leaf tissues of banana cv. Berangan ramets. (Values are means of four replicates. Vertical Bars represent Standard Error of Means)	4.35
4.2 Population of UPM31P1 isolated from root, corm, pseudostem and leaf tissues of banana cv. Berangan ramets. (Values are means of four replicates. Vertical Bars represent Standard Error of Means)	4.35

4.3	Population of UPM14B1 isolated from root, corm, pseudostem and leaf tissues of banana cv. Berangan ramets. (Values are means of four replicates. Vertical Bars represent Standard Error of Means)	4.36
4.4	Population of UPM13B8 isolated from root, corm, pseudostem and leaf tissues of banana cv. Berangan ramets. (Values are means of four replicates. Vertical Bars represent Standard Error of Means)	4.36
4.5	Population of UPM39B3 isolated from root, corm, pseudostem and leaf tissues of banana cv. Berangan ramets. (Values are means of four replicates. Vertical Bars represent Standard Error of Means)	4.37
5.1	Disease incidence recorded in ramets pre-inoculated with endophytes, according to days after challenged inoculation with FocR4. (Values were means of eight replicates)	5.13
5.2	Disease severity recorded in ramets pre-inoculated with endophytes, according to days after challenged inoculation with FocR4. (Values were means of eight replicates)	5.14
5.3	Disease incidence for ramets pre-inoculated with UPM31P1 and UPM31P1 + UPM39B3 at 16 weeks after challenged inoculation with FocR4. (Values are means of 25 replicates)	5.47
6.1	Percentage of total number of surviving plants pre-inoculated with UPM31P1 (TA) and UPM31P1 + UPM39B3 (TB), as compared to control (TC) observed up to 28 weeks in the field ('hot-spot'). (Values are means of 24 replicates). ( indicates foliar symptoms in Plate 6.6 to 6.8). ( indicates the first foliar symptom detected- 8 week for TC, week 14 for TA and TB)	6.9
6.2	Means of plant height (A), pseudostem diameter (B) and total number of leaves plant ⁻¹ (C) for plants pre-inoculated with UPM31P1 (TA) and UPM31P1 + UPM39B3 (TB), as compared to control (TC) at 12 weeks in the 'hot-spot'. (Values are means of 24 replicates. Vertical Bars represent Standard Error of Means)	6.16

6.3 Population of UPM31P1 (*F. oxysporum*) and UPM39B3 (*S. marcescens*) derived from root tissues and rhizosphere of plants pre-inoculated with UPM31P1 singly (TA) (A) and mixtures of UPM31P1 + UPM39B3 (TB) (B) after 1st and 2nd endophyte inoculation. (Values are means of seven replicates. Vertical Bars represent Standard Error of Means)

6.18

LIST OF PLATES

Plate	Page
3.1 The distribution of the sampling areas (1A); and the habitats of wild bananas selected for sampling (1B-1D)	3.3
3.2 Dual culture test: Preliminary <i>in vitro</i> screening for antagonistic activity of bacterial endophytes towards FocR4	3.7
3.3 Intracellular growth and colonization of host tissues by hyphae of fungal endophyte (→) as observed in root tissues of wild bananas (X400)	3.21
3.4 Scanning electron micrographs of normal structures of FocR4 hyphae in control plate (A) and abnormal hyphae structures upon exposure to bacterial (B and C) and fungal (D) endophytes recovered from inhibition zone in the dual culture test	3.29
3.5 Clear and reproducible DNA banding patterns of fungal isolates (isolate 1,2,3,4,5) produced by PCR product amplified with OPC primers 11, 14 and 15. (OPC 11, 14 and 15 were selected for subsequent RAPD-PCR analysis). (M: molecular marker 100 bp DNA Ladder Plus)	3.32
3.6 Clear and reproducible DNA banding patterns for bacterial isolates (isolate 1,2,3,4,5) produced by PCR product amplified with OPC primers 1, 8 and 11. (OPC 1, 8 and 11 were selected for subsequent RAPD-PCR analysis). (M: molecular marker 100 bp DNA Ladder Plus)	3.36
3.7 Cultural and morphological characteristics of 10 day-old cultures of UPM31F4 (A, B and C) and UPM 31P1 (D, E and F)	3.40
3.8 Colony pigmentation on NA and SEM micrograph of UPM14B1 (A and B), UPM 13B8 (C and D) and UPM39B3 (E and F)	3.42
4.1 Pictorial representation of the disease rating scale from 0 to 5 (A to F), based on extent of foliar symptoms in susceptible banana (Berangan cv. Intan) as used in disease severity assessment	4.6



4.2	Measurement of height (A) and pseudostem diameter (B) of banana (Berangan cv. Intan)	4.12
4.3	Respective parts of ramet (A) used for assessing the colonization ability of endophytes; roots (i), corm (ii), pseudostem (iii) and leaf (3 rd leaf) (iv). Roots were further divided to three different sections (B); roots from the base of the corm (Bs), from the middle region (Mid) and the root tips (Rt)	4.14
4.4	Reisolation of bacterial (A and B) and fungal (C and D) endophytes from tissue samples 24 hr after inoculation. (Bs-root base, Mid-middle region of root, Rt-root tips, corm, P-pseudostem, 3 rd leaf).	4.32
5.1	A: Development of Fusarium wilt in banana ramets pre-inoculated with UPM31P1 (T6) (ii), UPM31P1+UPM39B3 (T7) (iii), UPM39B3 (T8) (iv), UPM14B1+UPM13B8+UPM39B3 (T9) (v), at 28 days after challenged inoculation with FocR4. Control ramets from T5 (SDW-FocR4) and T10 (+FocR4) are shown in (i) and (vi), respectively. B: No reddish streaks or vascular discolouration was observed in uninfected control ramets, as compared to typical reddish streaks in both corm and pseudostem of infected ramets (C)	5.11
5.2	A: SEM micrograph of root tissues from treatment T4 (UPM14B1 + UPM13B8 + UPM39B3) showing the presence of bacteria (b) in parenchyma cells (p) of banana root tissues. B: Light Microscopy (100x) of endophytic colonization of UPM31P1 (\) spreads from xylem vessels (Xy) to adjacent angular parenchyma (AP) cells, but not to phloem cells (Ph) as observed in root tissues from T1 (UPM31P1). (Cell walls colonized by bacterial and fungal endophytes remained turgid and intact)	5.21
5.3	A: SEM micrograph of root tissues from treatment T10 (FocR4 only) showing the disintegration of xylem (Xy) vessels in banana root tissues. B: Light Microscopy (100x) showing disintegration of xylem (Xy) vessels (\) in root tissues of ramets from T10 (FocR4 only). Infected tissues appeared 'brittle' and lost its turgour when compared to control (C). (p-parenchyma tissues, Ph-phloem tissues)	5.22
5.4	Pigmentation of bacterial streaks on NA (left) and Cetrimide Agar (right) for UPM14B1 (A), UPM13B8 (B) and UPM39B3 (C)	5.41



5.5	Development of foliar symptoms of ramets in T2 (UPM31P1 + +FocR4) (A), T4 (UPM31P1 + UPM39B3 + FocR4) (B) and T6 (FocR4) (C) at 12 weeks after challenged with FocR4	5.48
6.1	Site for field testing in Field 53 (Block 8), Jenderata Estate, United Plantations Bhd., Teluk Intan, Perak. Transplanting of 8-week old ramets to Field 53 (A) and ramets in the field at four weeks after transplanting (B)	6.3
6.2	Inoculum used for field testing; UPM31P1 in 300g bags (i) and laminar air-dried inoculum as solid substrate (ii), and bacterial (UPM39B3) suspension in LB broth (iii)	6.3
6.3	Four-week old ramets treated with endophytes (A) and without endophytes (control) (B)	6.5
6.4	Application of UPM31P1 in substrate form (▼) (A) and bacterial suspension (UPM39B3) (B) into planting hole prior to planting of banana ramets	6.5
6.5	Plants from TA (UPM31P1) with no foliar symptoms (A) as compared to plants from TC (control) with yellowish streaks on the leaves (B) and reddish streaks in the pseudostem (C) and corm (D) tissues. (Plants were eight weeks in the field)	6.8
6.6	Development of foliar associated symptoms on plants with TA (UPM31P1) (A, B), TB (UPM31P1+ UPM39B3) (C, D) and TC (control) (E, F) at 12 weeks in the field ('hot-spot')	6.11
6.7	Development of foliar associated symptoms on plants with TA (UPM31P1) (A, B), TB (UPM31P1 + UPM39B3) (C, D) and TC (control) (E, F) at 16 weeks in the field ('hot-spot')	6.12
6.8	Development of foliar associated symptoms on plants with TA (UPM31P1) (A, B), TB (UPM31P1 + UPM39B3) (C, D) and TC (control) (E, F) at 20 weeks in the field ('hot-spot')	6.13



LIST OF ABBREVIATIONS

ANOVA	Analysis Of Variance
AP-PCR	Arbitrarily Primed-Polymerase Chain Reaction
AUDPC	Area Under Disease Progress Curve
BCA	Biological Control Agent
bp	Base Pair
BUG	Biolog Universal Media
c.f.u.	Colony Forming Units
CRD	Complete Randomized Design
DAF	DNA Amplification Fingerprinting
DI	Disease Incidence
DNA	Deoxyribonucleic Acid
dNTPs	Deoxyribonucleoside Triphosphates
DOA	Department of Agriculture
DR	Disease Reduction
DS	Disease Severity
EDTA	Ethylenediaminetetraacetic Acid
EMBRAPA	Brazilian Agricultural Research Cooperation
EMs	Endophytic Microorganisms
FAA	Formalin Acetic Acid
FAO	Food and Agriculture Organization
FHIA	Honduras Foundation of Agricultural Research